



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

PROBLEMS AND SOLUTIONS.

EDITED BY B. F. FINKEL AND OTTO DUNKEL.

Send all communications about Problems and Solutions to **B. F. FINKEL**, Springfield, Mo.

PROBLEMS FOR SOLUTION.

[N.B. The editorial work of this department would be greatly facilitated if, on sending in problems, the proposers would also enclose their solutions—*when they have them*. If a problem proposed is not original the proposer is requested *invariably* to state the fact and to give an exact reference to the source.]

2863. Proposed by A. A. BENNETT, Baltimore, Md.

From their generation as roulette curves, show that the two hypocycloids of five cusps drawn with common vertices, are such that each is the envelope of a chord of constant length suitably placed upon the other.

Show that for any odd prime p , the $(p - 1)/2$ distinct p -cusped hypocycloids with common vertices may be arranged in cycles, so that each is the envelope of a chord of constant length taken upon the succeeding curve of the cycle.

2864. Proposed by C. B. HALDEMAN, Ross, O.

If S is a side of a regular undecagon inscribed in a circle of radius unity, show that

$$S^5 - (S^4 - 3S^2 - 1)\sqrt{11} - 11S = 0.$$

2865. Proposed by JOSEPH ROSENBAUM, Milford, Conn.

In a circle, a chord AB is fixed in position and a moving chord CD is constant in length. Find the locus of the intersection of the bisectors of the angles ACD and BDC .

2866. Proposed by NORMAN ANNING, University of Michigan.

Equilateral triangles whose sides are 1, 3, 5, 7, ... are placed so that their bases lie corner to corner in a straight line. Show that the vertices lie upon a parabola and are all at integral distances from its focus.

2867. Proposed by W. H. HAYS, Colton, Wash.

Show that the probable value of the sum of the squared differences, got by pairing two sets of n numbers each, at random, is $n(n^2 - 1)/6$; and that the coefficient of correlation in such a case is 0. See Thorndike, *Mental and Social Measurements*, page 164.

2868. Proposed by H. S. UHLER, Yale University.

Let the evolute of a given curve be called the evolute of the first order, let the evolute of the first evolute be called the evolute of the second order, etc. Then, being given the following parametric equations in which a is a constant and γ is the parameter, namely,

$$x = (1 + 2 \sin^2 \gamma) \cos \gamma - a \sin 2\gamma,$$

$$y = 2 \sin^3 \gamma + a \cos 2\gamma$$

- find: (a) the parametric equations of the evolute of order n , both for n even and for n odd;
 (b) a formula for the total length of the n th evolute;
 (c) a formula for the total area of the n th evolute;
 (d) the sum of the lengths of all the evolutes from; $n = 1$ to $n = \infty$; and
 (e) the sum of the areas of all the evolutes from $n = 1$ to $n = \infty$.

Note: The original equations represent the envelope required in problem 2819, 1920, 134.